CLOZASOL DRENCH کلوزاسول ڈرینچ

Levamisole HCl + Oxyclozanide

لیوامیسول ہائیڈروکلورائیڈ + اوکسی کلوزانائیڈ

A Broad spectrum anthelmintic with the combination of two anthelmintics with alternative modes of action. For the treatment and control of both gastro-intestinal and pulmonary nematode infections, and adult liver fluke infections in cattle, sheep and goats.

ایک وسیع العمل انتہائی موثر کرم کُش دوا ہے جو چھوٹے بڑے جانوروں میں آنت، جگر، معدہ اور پھیپھڑوں میں موجود بالغ و نابالغ گول کرم، ٹیپ ورم اور فیشولا کا خاتمہ کرتی ہے۔



Manufactured by:

WIMITS

Wimits Pharmaceuticals (Pvt.) Ltd.

Empowering animal health

CLOZASOL DRENCH

COMPOSITION:

Each 100 ml contains
Levamisole HCl (BP).....1.50 gm
Oxyclozanide (BP Vet.).....3.00 gm

INDICATIONS:

Clozasol drench is a broad spectrum anthelmintic for use in the treatment and control of both gastro-intestinal and pulmonary nematode infections and adult liver fluke infections in cattle, sheep and goats. It should be used in cases of parasitic gastroenteritis and lungworm disease caused by mature and developing immature forms of those organisms sensitive to treatment with levamisole hydrochloride.
Lungworms: Dictyocaulus spp.

Gastrointestinal worms: Haemonchus spp; Ostertagia spp (except inhibited Ostertagia larvae in cattle); Nematodirus spp; Trichostrongylus spp; Cooperia spp; Oesophagostomum spp; and Bunostomum spp.

Clozasol drench also removes most mature Fasciola spp; (flukes) present in the bile ducts of the liver. It is a single dose treatment for cattle, sheep and goats effective against all major stomach and bowel worm species (including Ostertagia and Nematodirus), Lungworms (causing 'Husk' or "Hoose"), Tapeworm segments and Liver flukes.

Clozasol drench due to its wide safety margins is suitable for stock of all ages, including young, pregnant and lactating animals. No diet restrictions. Animal need not to be fasted, taken off concentrates or test dosed.

DOSAGE:

1 ml per 3 Kg B.W.
Clozasol drench should be administered as an oral drench using a suitable drenching gun.

WITHDRAWAL TIMES

Meat : 28 days. Milk : 4 days.

PRECAUTIONS:

Protect from heat and sunlight.
Store in a cool & dry place at 25°C.
Keep out of reach of children.
Consult veterinarian before use.
Harmful if absorbed by skin contact or swallowed by humans. Wash hands after use.
Shake well before use.

PACKING:

100 ml and 1 litre.

گول کرموں کے انڈوں اور لاروا کو بھی تلف کرتی ہے۔

خوراک واستعمال:

املی لیٹر بحساب ۳ کلو جسمانی وزن چھوٹے بڑے جانوروں میں بذریعہ ڈرینچنگ گن پلائیں۔

احتیاط:

گرمی اور دھوپ سے بچائیں۔ تمام دوائیں بچوں کی پہنچ سے دور رکھیں۔ ڈاکٹر کی ہدایات کے مطابق استعمال کریں۔ جلد پر لگنے یا نگلنے کی صورت میں نقصان دہ ہے۔ استعمال کے بعد ہاتھوں کو دھولیں۔ استعمال سے پہلے بوتل اچھی طرح ہلالیں۔

اجزائے ترکیبی:

فی ۱۰۰ املی لیٹر

لیوامیسول ہائیڈروکلورائیڈ (بی پی)۔۔۔۔۔۔۔۔۔۔۔۔۔۔۔۔۔۱ء۵ گرام

اوکسی کلوزانائیڈ (بی پی۔ویٹ)۔۔۔۔۔۔۔۔۔۔۔۔۔۔۔۔۔