

# Donits

(Domperidone)  
Suspension

**WIMITS**  
Pharmaceuticals (Pvt) Ltd.

Donits used to treat nausea and vomiting, relief from symptoms associated with gastroparesis including loss of appetite, bloating & belching with and after meal.

**Composition:** Each ml contains Domperidone 1 mg

**Pharmacological Group:** Anti-emetic

## Indications

- It is indicated in the treatment of epigastric sense of fullness.
- Upper abdominal distension
- Regurgitation
- Nausea and Vomiting

**Proposed Route of Administration:** Oral liquid Suspension  
It is recommended to take oral Domperidone before meals. If taken after meals, absorption of the drug is somewhat delayed.

## Proposed Dosage:

Actual Dosage must be determined by the Physician:

## Adults:

### Adults and adolescents (over 12 years and weighing 35 kg or more)

The initial duration of treatment is four weeks. Patients should be re-evaluated after four weeks and the need for continued treatment re-assessed. 10 ml to 20 ml (of oral suspension containing Domperidone 1mg per ml) three to four times per day with a maximum daily dose of 80 ml.

## Infants and children

0.25 – 0.5 mg/kg three to four times per day with a maximum daily dose of 2.4 mg/kg (but do not exceed 80 mg per day).

**Proposed Storage Conditions:** Store below 25°C in a dry and cool place. Protect from heat, light and moisture.

**Pack Size:** 60ml, 120ml suspension, Tablets



**WIMITS**  
Pharmaceuticals (Pvt) Ltd.

Manufactured By:  
**Wimits Pharmaceuticals (Pvt) Ltd.**  
129 – Sunder Industrial Estate Lahore, Pakistan  
UAN: 0092-42-111-11-6366, Fax: +92-42-35462418, Email: [info@wimits.com](mailto:info@wimits.com)

