



**Ebalin** is a medication used to treat epilepsy, neuropathic pain, and fibromyalgia.

**Composition:** Each capsule contains Pregabalin 150mg

Pharmacological Group: Anticonvulsant

## **Indications:**

- Management of neuropathic pain associated with diabetic peripheral neuropathy
- Management of post herpetic neuralgia
- Adjunctive therapy for adult patients with partial onset seizures
- Management of fibromyalgia
- Management of neuropathic pain associated with spinal cord injury



## **Mechanism of Action**

Pregabalin binds with high affinity to the alpha2-delta site (an auxiliary subunit of voltage-gated calcium channels) in central nervous system tissues.

## **Pharmacokinetics**

Pregabalin is well absorbed after oral administration, is eliminated largely by renal excretion, and has an elimination half-life of about 6 hours.

**Proposed Route of Administration:** Oral (Capsule)

## **Proposed Dosage:**

Pregabalin may be taken with or without food. The initial dose for neuropathic pain associated with diabetic peripheral neuropathy is 50 mg three times a day (150 mg/day). For other indication use as directed by qualified physician.

**Proposed Storage Conditions:** Store below 25°C in a dry and cool place. Protect from heat, light and moisture.

Pack Size: 14's capsules



Manufactured By:

Wimits Pharmaceuticals (Pvt) Ltd. 129 – Sunder Industrial Estate Lahore, Pakistan. UAN: 0092-42-111-11-6366,

Fax: +92-42-35462418 Email: info@wimits.com





