



Fenbid used to treat painful conditions such as Toothache, arthritis, sprains and strains, and migraine pain and pain after surgical operations.

Composition: Each film coated tablet contains Flurbiprofen BP 100mg

Pharmacological Group: Antipyretic, Analgesic, Anti-Inflammatory

Indication:

- Fenbid is also indicated for its analgesic effect in the relief of mild to moderate pain in conditions such as dental pain, post-operative pain, dysmenorrhea and migraine.
- Fenbid for the treatment of rheumatoid disease, osteoarthritis, ankylosing spondylitis, musculoskeletal disorders and trauma such as periarthritis, frozen shoulder, bursitis, Tendinitis, tenosynovitis, low back pain, sprains and strains.



Proposed Route of Administration: Oral (tablet) **Proposed Dosage:**

Actual Dosage must be determined by the Physician:

Adults: 100 to 200 mg daily in two, three or four divided doses. In patients with severe symptoms or disease of recent origin, or during acute exacerbations, the total daily dosage may be increased to 300 mg in divided doses.

Dysmenorrhoea: A dosage of 100 mg may be administered at the start of symptoms followed by 50 or 100 mg given at four- to six-hour intervals.

The maximum total daily dosage not to exceed 300 mg.

Children: Not recommended for use in children under 12 years.

Elderly: The elderly are at increased risk of the serious consequences of adverse reactions. If an NSAID is considered necessary, the lowest dose should be used and the patient should be monitored for GI bleeding for four weeks following initiation of NSAID therapy. Undesirable effects may be minimized by using the lowest effective dose for the shortest duration necessary to control symptoms.


Proposed Storage Conditions: Store below 25C in a dry and cool place. Protect from heat, light and moisture.

Pack size: 30 Tablets











Fax: +92-42-35462418,Email: info@wimits.com