

# LEVACLOZ BOLUS

## ليواكلوز بولس

Levamisole + Oxyclozanide



Manufactured by:

**WIMITS**

Wimits Pharmaceuticals (Pvt.) Ltd.  
Plot No 129 Sunder Industrial Estate Lahore.

Serving Animal Health With Pride

# LEVACLOZ BOLUS

## COMPOSITION:

Levamisole HCL(BP)...1125 mg/bolus  
Oxyclozanide (BP Vet.)..2250 mg/bolus

## INDICATIONS:

LEVACLOZ BOLUS is a broad spectrum anthelmintic for use in the treatment and control of both gastrointestinal and pulmonary nematode infections and adult liver fluke infections in cattle and sheep only. LEVACLOZ BOLUS should be used in cases of parasitic gastroenteritis and lungworm disease caused by mature and developing immature forms of those organisms sensitive to treatment with levamisole hydrochloride.

## Lungworms: Dictyocaulus spp.

**Gastrointestinal worms:** Haemonchus spp; Ostertagia spp (except inhibited Ostertagia larvae in cattle); Nematodirus spp; Trichostrongylus spp; Cooperia spp; Oesophagostomum spp; and Bunostomum spp.

It also removes most mature Fasciola spp; (flukes) present in the bile ducts of the liver. It is a single dose treatment for cattle, sheep and goats effective against; All major stomach and bowel worm species (including Ostertagia and Nematodirus). Lungworms (causing 'Husk' or 'Hoose'). Tapeworm segments. Liver flukes.

LEVACLOZ BOLUS due to its wide safety margins is suitable for stock of all ages, including young, pregnant and lactating animals. No diet restrictions. Animal need not to be fasted, taken off concentrates or test dosed.

## CONTRAINDICATIONS, WARNINGS, Etc:

To ensure administration of a correct dose, bodyweight should be determined as accurately as possible; accuracy of the dosing device should be checked.

Under dosing, which may be due to under estimation of body weight, mis administration of the product, or lack of calibration of the dosing device.

## DOSAGE & ADMINISTRATION:

1 Bolus per 175kg body weight.

LEVACLOZ BOLUS should be administered as an oral drench. Dosing must be carried out accurately using a suitable gun system.

## PRECAUTIONS:

### PROTECTION OF OPERATORS:

No special precautions are necessary for the protection of operators, when treating cattle, sheep and goats with Zelcon Plus.

### PROTECTION OF LIVESTOCK, WILDLIFE AND OTHERS:

The use of LEVACLOZ BOLUS does not present a hazard to livestock, wildlife and others. Wash out container thoroughly and dispose off safely.

### WITHDRAWAL PERIOD:

Animals must not be slaughtered for human consumption during treatment. Cattle and sheep may be slaughtered for human consumption only after 14 days from the last treatment.

Milk taken from lactating animals in the 24 hours following treatment should not be used for human consumption.

## PRECAUTIONS:

Store in a dry, dark place between 2°C and 25°C.

Store in the well closed blister packs. Keep medicine away from children.

To be used as directed by the registered veterinary practitioners only.

**PACKING:** Pack of 1 x 50

Serving Animal Health With Pride