

# LOXITAR

Injection

انجیکشن  
لوکسی ٹار

Alleviation of fever, inflammation and pain in both acute and chronic musculo-skeletal disorders.

جانوروں میں سوزش، درد اور بخار کی موثر دوا

Meloxicam Injection



میلو کسیمی انجیکشن



Manufactured by:

**WIMITS** Wimits Pharmaceuticals (Pvt.) Ltd.

# LOXITAR لوکسی ٹار

Injection

انجیکشن

## COMPOSITION

Each ml contains  
Meloxicam.....10mg

## INDICATIONS

Meloxicam is a non-steroidal anti-inflammatory drug (NSAID) derived from oxicam. Meloxicam acts by inhibition of prostaglandin synthesis exerting anti-inflammatory, anti-exudative, analgesic and antipyretic effects. It reduces leukocyte infiltration into the inflamed tissue.

Loxitar injection is highly effective against Fever of infective origins, Rheumatic Fever, Arthritis, Polyarthritis, All types of Surgical, Traumatic & Infective inflammatory & Painful conditions. For use in acute respiratory infection with appropriate antibiotic therapy to reduce clinical signs in cattle. For use in diarrhoea in combination with oral re-hydration therapy to reduce clinical signs in calves of over one week of age and young, non-lactating cattle. For adjunctive therapy in the treatment of acute mastitis, in combination with antibiotic therapy.

## DOSAGE AND ADMINISTRATION

Inject by intramuscular route only.  
Horses and Camels.....7.5-11.25 ml  
Cows and Buffaloes.....06-7.5 ml  
Sheep and Goats.....1.5-2.25 ml

The recommended maximum volume to be administered at a single injection site is 10ml. A suitable oral therapy containing meloxicam, may be used for continuation of treatment. Avoid introduction of contamination during use.

اونٹ و گھوڑے.....۷.۵-۱۱.۲۵ ملی لیٹر  
گائے و بھینس.....۶-۷.۵ ملی لیٹر  
بھینر و بکری.....۱.۵-۲.۲۵ ملی لیٹر

احتیاط:

گرمی اور دھوپ سے بچائیں۔ ٹھنڈی اور خشک جگہ پر ۲۵ ڈگری سینٹی گریڈ درجہ حرارت پر رکھیں۔ تمام دوائیں بچوں کی پہنچ سے دور رکھیں۔ وینٹری ڈاکٹر کی ہدایات کے مطابق استعمال کریں۔ سبیل کے ٹیک ہونے، دھندلا ہونے یا اس میں کوئی غیر حل پذیر شے نظر آنے کی صورت میں ہرگز استعمال نہ کریں۔

## CONTRAINDICATIONS, WARNINGS, Etc

Do not use in horses less than 6 weeks of age. Do not use in pregnant or lactating mares. Do not use in animals suffering from impaired hepatic, cardiac or renal function and haemorrhagic disorders, or where there is evidence of ulcerogenic gastrointestinal lesions.

Do not use in case of hypersensitivity to the active substance or to any of the excipients. For the treatment of diarrhoea in cattle. Do not use in animals of less than one week of age. If adverse reactions occur, treatment should be discontinued and the advice of a veterinarian should be sought. Avoid use in very severely dehydrated, hypovolaemic or hypotensive animals which require parenteral rehydration, as there may be a potential risk of renal toxicity.

In case of inadequate relief of pain when used in the treatment of equine colic, careful re-evaluation of the diagnosis should be made as this could indicate the need for surgical intervention.

## USE DURING PREGNANCY AND LACTATION

Can be used during pregnancy and lactation in cattle. Do not administer concurrently with gluco-corticosteroids, other non-steroidal anti-inflammatory drugs or with anticoagulant agents. In the case of overdose, symptomatic treatment should be initiated.

## PRECAUTIONS

Protect from moisture, freezing, heat & sunlight. Keep all medicines out of the reach of children. To be used as directed by the registered veterinary practitioners only. Don't use if the seal is leaking, solution is turbid or contains any foreign matter.

اجزائے ترکیبی:

نی ملی لیٹر

میلو کسیمی (بی بی).....۱۰ ملی گرام

علامت:

لوکسی ٹار انجیکشن جانوروں میں درد، سوزش، جوڑوں کے درد، بخار اور سوزی تھکاوٹ کے لیے موثر دوا ہے۔

خوراک و استعمال:

جانوروں میں درج ذیل خوراک، زیر جلد یا گوشت میں انجیکشن لگائیں

129-Sunder Industrial Estate Lahore.

info@wimits.com

Phone: 042-3750291, Fax: 042-3750292,

www.wimits.com