

Macral is macrolide antibiotic used to treat bacterial infections such as respiratory tract infections, ear and skin infections and is used in combination with other antibiotics to eradicate H.pylori in duodenal ulcers.

Composition: Each film coated tablet contains: Clarithromycin BP 250mg, 500mg

Pharmacological Group: Macrolide

- Bacterial pharyngitis
- Mild to moderate community-acquired pneumonia
- Acute bacterial sinusitis (adequately diagnosed)
- Acute exacerbation of chronic bronchitis
- Skin and soft tissue infections of mild to moderate severity.



In appropriate combination with antibacterial therapeutic regimens and an appropriate ulcer healing medicinal product for the eradication of Helicobacter pylori in adult patients with Helicobacter pylori-associated ulcers.

Proposed Dosage:

The dosage of Clarithromycin film-coated tablets depends on the type and severity of the infection and has to be defined in any case by the physician.

Adults:

Standard dosage: The usual dose is 250mg twice daily (in the morning and in the evening)

High dosage treatment (severe infections): The usual dose may be increased to 500 mg twice daily in severe infections. Children older than 12 years: As for adults.

Children younger than 12 years: Use of Clarithromycin 250mg Tablets are not recommended for children younger than 12 years. Use clarithromycin Paediatric Suspension.

Proposed Route of Administration: Oral (tablet)

Proposed Storage Conditions: Store below 30°C in a dry and cool place. Protect from heat, light and moisture.

Pack size: 10 Film coated tablets of 250mg, 500mg.