

For I.M. / I.V. Use

MECOWIM
Mecobalamin 500mcg

WIMITS
Pharmaceuticals (Pvt) Ltd.

Mecowim injection is prescribed for various types of nerve pain, peripheral neuropathy, megaloblastic anaemia, and as a preliminary treatment for amyotrophic lateral sclerosis

Composition: Each ml contains: Mecobalamin.500mcg.

Pharmacological Group: vitamin, hematinic

Indications:

Mecowim is indicated for peripheral neuropathies, Pernicious anemia and megaloblastic anemia due to vitamin B12 deficiency.

Proposed Route of Administration: Intra muscular (IM) Injection or Intra venous (IV) Injection

Proposed Dosage:

Peripheral neuropathies:

Adults: IM or IV injection three times a week

Megaloblastic anaemia:

Adults:

IM or IV injection three times a week. After two months 1 ampoule in a single administration at 1 to 3 months intervals for maintenance therapy

Proposed Storage Conditions: Store below 25°C. Protect from light and heat.

Adverse Reactions/Side Effects

Mecowim is nontoxic, even in large doses. Adverse reactions reported following administration include headache, infection, nausea/vomiting, parenthesis, and rhinitis.

Pack Size: 10 x 1ml ampoules, Tablets 20, 50, 100



Manufactured By:

WIMITS
Pharmaceuticals (Pvt) Ltd.

Wimits Pharmaceuticals (Pvt) Ltd.
129 – Sunder Industrial Estate Lahore, Pakistan
UAN: 0092-42-111-11-6366,
Fax: +92-42-35462418, Email: info@wimits.com

