

Iron Syrup

P-MALT

Iron (III)-hydroxide
Polymaltose Complex

WIMITS
Pharmaceuticals (Pvt) Ltd.

Composition: Each 5ml contains Iron III Hydroxide Polymaltose Complex.. 50 mg

Pharmacological Group: Anti-Anaemics, Haematinic

P-Malt indicated for the treatment of latent iron deficiency and iron deficiency anemia during and after pregnancy (during lactation). For prophylactic therapy of iron deficiency to cover the recommended daily dietary allowances.

Proposed Route of Administration: Oral liquid

Proposed Dosage:

Actual Dosage must be determined by the Physician:

Adults: P-Malt dosage and duration of therapy is dependent upon iron deficiency. The daily dose can be taken at one time or it can be divided into separate doses. In case of manifested iron deficiency, the therapy takes about 3-5 months until normalization of hemoglobin value is achieved. Afterward the therapy should be continued for several weeks with a dosage such as described for latent iron deficiency to replenish the iron stores.

Children: 6 – 12 Years: 5 ml daily

Above 12 years, Adults and nursing women:

- Manifest Iron Deficiency: 5 – 15 ml daily
- Latent Iron Deficiency: 5 ml daily

During Pregnancy:

- Manifest Iron Deficiency: 5 – 15 ml daily
- Latent Iron Deficiency: 5 ml daily

Prophylactic Therapy: 2.5 ml daily

Proposed Storage Conditions: Store below 25C in a dry and cool place. Protect from heat, light and moisture.

Pack Size: 60 ml, 120ml syrup



WIMITS
Pharmaceuticals (Pvt) Ltd.

Manufactured By:
Wimits Pharmaceuticals (Pvt) Ltd.
129 – Sunder Industrial Estate Lahore, Pakistan
UAN: 0092-42-111-11-6366, Fax: +92-42-35462418
Email: info@wimits.com

