

TRICLAMEX

(Triclabendazole + Levamisole HCl)

Oral combination of a Broad Spectrum de-wormer and a wholly effective flukicide

ایک وسیع العمل کرم کش اور فلوک کش دوا

ڈرینج ٹرائیکلامیکس

ٹرائیکلابنڈازول + لیوامیسول ہائیڈروکلورائیڈ



Manufactured by:

WIMITS Pharmaceuticals (Pvt.) Ltd.
Empowering animal health

TRICLAMEX DRENCH

ڈرینج ٹرائیکلامیکس

COMPOSITION:

Each ml Contains
Triclabendazole50 mg
Levamisole37.50 mg


INDICATIONS:

For the treatment and control of parasitic bronchitis, parasitic gastro-enteritis and fascioliasis in cattle. When used at the recommended dose rate, the product is effective against mature and developing immature stages of Levamisole susceptible Haemonchus, Trichostrongylus, Cooperia, Nematodirus, Bunostromum and Oesophagostomum species in the gastro-intestinal tract, and Dictyocaulus spp in the lungs. The product is effective against developing immature and adult Ostertagia, but is not active against inhibited larvae.

It is also effective against all stages of Triclabendazole susceptible Fasciola hepatica from two week old early immature to adult fluke, and so is indicated against both acute and chronic fascioliasis.

Triclamex drench contains Levamisole which is a broad spectrum anthelmintic that is effective against mature and immature stages of nematodes, including strains resistant to other anthelmintics.

Triclabendazole is specifically effective against all 3 stages of Fasciola. The unique efficacy against early mature and immature stages makes Triclamex drench particularly useful for the treatment of outbreaks of acute, subacute and chronic fasciolosis and for the strategic control of Fasciola spp. infections, fascioliasis (either F. hepatica or F. gigantica) affecting cattle,



buffaloes, sheep and goats. Triclabendazole is a member of the benzimidazole family of drenches, but unlike the other members it is liver fluke specific. It is by far the most effective fluke molecule on the market, giving better control than any other flukicidal active.

Adult and immature forms of round worms such as: Haemonchus, Oestertagia, Trichostrongylus, Cooperia, Bunostomum, Nematodirus and Oesophagostomum affecting cattle, buffaloes, sheep and goats.

DOSAGE & ADMINISTRATION:

To be given as a oral drench at the rate of 1ml per 5 Kg body weight by drenching gun.

PRECAUTIONS:

Store below 30°C (room temperature) in original container tightly closed, in a cool dry place. Keep away from children. Consult Veterinarian before use. Assess body weight as accurately as possible before calculating the dosage. Clean drenching equipment before and after use. Use undiluted product from the original container. Intensive use or misuse of anthelmintics can give rise to resistance.

To reduce this risk, dosing programmes should be discussed with your Veterinarian.

Shake well before use.

PACKING:

100 ml and 1 litre.



خوراک و استعمال:

چھوٹے، بڑے جانوروں میں املی لیٹر فی ۵ کلوگرام بحساب جسمانی وزن بڑیہ ڈرینجنگٹ گن پائیں۔

احتیاط:

گرمی اور دھوپ سے بچائیں۔ تمام دواہیں بچوں کی پہنچ سے دور رکھیں ڈاکٹر کی ہدایات کے مطابق استعمال کریں۔ جلد پر لگنے یا لٹکنے کی صورت میں نقصان دہ ہے۔ استعمال کے بعد ہاتھوں کو دھو لیں۔

استعمال سے پہلے بوتل اچھی طرح ہلائیں۔

اجزائے ترکیبی:

فی ملی لیٹر

ٹرائیکلابنڈازول.....۵۰ ملی گرام

لیوامیسول ہائیڈروکلورائیڈ (بی بی).....۳۷.۵۰ ملی گرام

علامات:

ٹرائیکلامیکس ڈرینج میں موجود لیوامیسول اور ٹرائیکلابنڈازول ایکٹ دوسرے کی طاقت کو بڑھاتے ہیں جس سے جانوروں کے معدے میں موجود گول کرم، چکرے کرم، ٹیپ ورم، لیور فلوک اور پھیپھڑوں کے تمام بالغ اور نابالغ اقسام کا خاتمہ ہوتا ہے۔

129-Sunder Industrial Estate Lahore.

info@wimits.com

Phone: 042-37502921, Fax: 042-37502922,

www.wimits.com