

Each Film coated Tablet contains Desloratadine..... 5mg

Pharmacological Group: Antihistamine

Recommended Clinical Use: along with documentary evidence of the approved indications in the country of origin & in developed countries.

Indications: Seasonal Allergic Rhinitis, Perennial Allergic Rhinitis, Chronic Idiopathic Urticaria:

Desotin is indicated for the relief of the nasal and nonnasal symptoms of seasonal allergic rhinitis in patients 2 years of age and older.

Desotin is indicated for the symptomatic relief of pruritus, reduction in the number of hives, and size of hives, in patients with chronic idiopathic urticaria 6 months of age and older.

Mechanism of Action: Desloratadine is a long-acting tricyclic histamine antagonist with selective H₁-receptor histamine antagonist activity. .

Proposed Route of Administration: Oral (Tablet)

Dosage:

Adults and children 12 years of age and over:

5 mg tablet once daily OR as directed by Physician

Children 6 to 11 years of age:

Dose of Desotin Syrup is 1 teaspoonful (2.5 mg in 5 mL) once daily or the recommended dose of desloratadine Tablets is one 2.5 mg tablet once daily.

- **Children 12 months to 5 years of age:**
- Of desloratadine Syrup is 1/2 teaspoonful (1.25 mg in 2.5 mL) once daily.
- **Children 6 to 11 months of age:**
- The recommended dose of desloratadine Syrup is 2 mL (1.0 mg) once daily. The age-appropriate dose of desloratadine Syrup should be administered with a commercially available measuring dropper or syringe that is calibrated to deliver 2 mL and 2.5 mL (1/2 teaspoon).

In adult patients with liver or renal impairment, a starting dose of one 5 mg tablet every other day is recommended based on pharmacokinetic data. Dosing recommendation for children with liver or renal impairment cannot be made due to lack of data.

Storage Conditions: Store below 25C in a dry and cool place. Protect from heat, light and moisture.

Pack Size: 10's

