



K-lac a nonsteroidal anti-inflammatory drug (NSAID), is indicated for the short-term (up to 5 days in adults), management of moderately severe acute pain.

Composition: Each ml contains Ketorolac Trometamine... 30mg

Pharmacological Group: Cyclo-Oxygenase Inhibitor, Analgesic, Anti-Inflammatory.

Indications:

Short-term management of moderate to severe post-operative pain.

Proposed Route of Administration: Intra muscular (IM) Injection or Intra venous (IV) Injection

Proposed Dosage:

Adults: 16 years to <65 years: single dose: 60 mg IM or 30 mg IV.

Multiple dose: 30 mg IV/IM every 6 hours. Max: 120mg/day for 2 days.

aays.

>65 years or <50 kg: single dose: 30 mg IM or 50 mg IV.

Multiple doses: 15 mg IV/IM every 6 hours. Max: 60 mg/day for 2 days.

Children: Under 16 years, not recommended.

Proposed Storage Conditions: Store below 30°C. Protect from light and heat.

Pack size: 1*5Ampoules







Manufactured By:



Wimits Pharmaceuticals (Pvt) Ltd. 129 – Sunder Industrial Estate Lahore, Pakistan UAN: 0092-42-111-11-6366, Fax: +92-42-35462418

Email: info@wimits.com

