

Solo-Cip

250mg 500mg
(Ciprofloxacin)

Solo-cip is an antibiotic used to treat bacterial infections includes intra-abdominal infections, diarrhea, respiratory tract infections, skin infections, typhoid fever, and urinary tract infections.

Composition: Each Film Coated Tablet contains Ciprofloxacin as Hydrochloride 250mg / 500mg

Pharmacological Group: Antibiotic, Quinolone

Indications:

- Acute maxillary sinusitis, bacterial exacerbation of chronic bronchitis
- Community-acquired pneumonia and nosocomial Pneumonia, Typhoid & paratyphoid fever
- Complicated & Uncomplicated skin and skin structure infections, wound infections
- Complicated & Uncomplicated UTI,



DOSAGE GUIDELINES

Infection	Severity	Dose	Frequency	Usual Durations*
Urinary Tract	Acute Uncomplicated	250 mg	q 12 h	3 days
	Mild/Moderate	250 mg	q 12 h	7 to 14 days
	Severe/Complicated	500 mg	q 12 h	7 to 14 days
Chronic Bacterial Prostatitis	Mild/Moderate	500 mg	q 12 h	28 days
Lower Respiratory Tract	Mild/Moderate	500 mg	q 12 h	7 to 14 days
	Severe/Complicated	750 mg	q 12 h	7 to 14 days
Acute Sinusitis	Mild/Moderate	500 mg	q 12 h	10 days
Skin and Skin Structure	Mild/Moderate	500 mg	q 12 h	7 to 14 days
Bone and Joint	Severe/Complicated	750 mg	q 12 h	7 to 14 days
	Mild/Moderate	500 mg	q 12 h	≥ 4 to 6 weeks
Intra-Abdominal†*	Severe/Complicated	750 mg	q 12 h	≥ 4 to 6 weeks
	Complicated	500 mg	q 12 h	7 to 14 days
Infectious Diarrhea	Mild/Moderate/Severe	500 mg	q 12 h	5 to 7 days
Typhoid Fever	Mild/Moderate	500 mg	q 12 h	10 days
Urethral and Cervical Gonococcal Infections	Uncomplicated	250 mg	single dose	single dose
Inhalational anthrax (post-exposure)‡		500 mg	q 12 h	60 days

Proposed storage conditions: Store below 25°C in a dry and cool place. Protect from heat, light and moisture.

Pack Size: 10 Film coated tables, 250mg, 500mg

Manufactured By: